



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 :

A61K 9/72, 47/26

A1

(11) International Publication Number:

WO 91/11179

(43) International Publication Date:

8 August 1991 (08.08.91)

(21) International Application Number: PCT/GB91/00103

(22) International Filing Date: 24 January 1991 (24.01.91)

(30) Priority data:
9001635.3

24 January 1990 (24.01.90) GB

(71) Applicant (for all designated States except US): NATIONAL
RESEARCH DEVELOPMENT CORPORATION
[GB/GB]; 101 Newington Causeway, London SE1 6BU
(GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GANDERTON, David
[GB/GB]; Crooked Chimneys, Cheriton Bishop, Exeter
EX6 6JL (GB). KASSEM, Nuha, Mohammed [Stateless/
GB]; 73 Latymer Court, Hammersmith Road, London
W6 7JE (GB).(74) Agent: HAMILTON, Raymond; Patents Department, Na-
tional Research Development Corporation, 101 Newing-
ton Causeway, London SE1 6BU (GB).(81) Designated States: AT, AT (European patent), AU, BB, BE
(European patent), BF (OAPI patent), BG, BJ (OAPI
patent), BR, CA, CF (OAPI patent), CG (OAPI patent),
CH, CH (European patent), CM (OAPI patent), DE,
DE (European patent), DK, DK (European patent), ES,
ES (European patent), FI, FR (European patent), GA
(OAPI patent), GB, GB (European patent), GR (Euro-
pean patent), HU, IT (European patent), JP, KP, KR,
LK, LU, LU (European patent), MC, MG, ML (OAPI
patent), MR (OAPI patent), MW, NL, NL (European
patent), NO, PL, RO, SD, SE, SE (European patent),
SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI
patent), US.

Published

With international search report.

(54) Title: AEROSOL CARRIERS

(57) Abstract

Pharmaceutical excipients useful in dry powder inhalents comprise particles having a rugosity (measured by air permeame-
try) of less than 1.75. The use of these carriers increases the amount of drug injected by the patient using a dry powder inhaler.
The preferred excipients are crystalline sugars such as lactose which may conveniently be prepared by controlled crystallisation
from an aqueous medium.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

- 1 -

AEROSOL CARRIERS

This invention relates to novel carrier materials useful in the formulation of pharmaceutical compositions especially dry powder compositions which are suitable for use in inhalation aerosols and to novel processes for the production of these materials.

The administration of pharmacological agents by inhalation has been recognised as a valuable technique, particularly in the treatment of diseases of the respiratory tract. The efficacy of the technique has been limited by difficulty in making appropriate dosages available to the lungs. The delivery systems currently available are nebulisers, pressurised metered dose inhalers and dry powder inhalers. Nebulisers are relatively effective but they are expensive and bulky and as a result are mainly used in hospitals. Pressurised metered dose inhalers require good co-ordination of actuation and inhalation which presents difficulties to many patients. They also require the use of propellants which may be undesirable on environmental grounds.

A variety of dry powder inhalers have been developed. All of them rely upon the inspiratory effort of the patient to produce finely divided drug particles which are available to the lungs. Also there have been various proposals for dry powder formulations suitable for use in these inhalers in order to improve the efficacy of the treatment. For example International Patent Application WO 87/05213 describes a carrier which comprises microgranules of a conglomerate of one or more solid water soluble diluents with a lubricant such as magnesium stearate. In practice another difficulty is caused by the tendency of the drug particles which are necessarily of a relatively small size to agglomerate either with themselves or more usually with particles of the carrier materials with which they are admixed. The difficulties inherent in redispersion of these agglomerates means that only a small proportion of the drug, may be as little as 5% is actually injected via the lungs.

- 2 -

The present invention is directed to novel materials which are useful as carriers in dry powder inhaler compositions. We have discovered that the redispersion of drug particles from compositions comprising carriers is facilitated if the rugosity of the carrier particles is reduced. The rugosity values of the materials are those measured by air permeametry. Accordingly, from one aspect our invention provides a particulate carrier suitable for use in the preparation of pharmaceutical compositions having an average particle size of from 5.0 to 1000 microns and a rugosity of less than 1.75. The measurement of rugosity by air permeametry produces a result which reflects the nature of the external surface of the material under test whereas measurements by techniques such as nitrogen adsorption reflect the total surface area including areas which are not accessible to particulate solids. The rugosity of conventional excipients measured by air permeametry has been found to be at least 1.96 and generally greater than 2.0. The carrier may be any crystalline non toxic material which is acceptable for use in pharmaceutical compositions which does not destabilise the pharmaceutically active materials with which it is formulated and which can be produced in a form having a rugosity of less than 1.75. The preferred carriers are those which are known to be useful in dry powder inhaler compositions especially the mono-saccharides such as lactose, mannitol, arabinose, xylitol and dextrose and their monohydrates, dissacharides such as maltose or sucrose and polysaccharides such as starches, dextrans or dextrans.

Preferably the carrier comprises a particulate crystalline sugar such as glucose, fructose, mannitol, sucrose and most preferably lactose.

The average size of the particles of the carrier is preferably in the range 5 to 1000 μm and more preferably in the range 30 to 250 μm and most preferably 50 to 100 μm . Typically at least 95% of the particles will be of a size which falls within this range, although the presence of significant

- 3 -

quantities of fine material may be tolerable albeit less preferred.

05 The particulate sugar crystals which constitute a preferred aspect, may be conveniently prepared by crystallisation from a solution which is preferably an aqueous solution. The conditions under which crystallisation occurs should be controlled so as to favour the production of crystals having the desired low degree of rugosity. In general conditions which allow the crystals to form slowly are preferred whilst those which result in rapid
10 crystallisation are correspondingly less preferred. The utility of any particularly crystallisation process must be evaluated empirically and it is within the skill of the art to modify unsatisfactory procedures in order to produce the desired crystalline form of the novel excipients.

15 Processes in which a sugar is precipitated from saturated aqueous solution by the addition of at least an equal volume of a water immiscible organic solvent and a quantity of a solvent which is miscible with both water and the aforesaid organic solvent which is at least 5% by volume of the total volume of the
20 aqueous solution and the organic solvent constitute another aspect of this invention. The novel precipitation process may be conveniently carried out by mixing the solution and the solvents at ambient temperature and maintaining them at that temperature with thorough mixing until sugar crystals are formed.

25 Seeding of the saturated solution may be advantageous insofar as it may reduce the time required for crystal formation.

The size and morphology of the particulate material may be varied by controlling the conditions under which crystallisation and crystal growth occurs. In particular, the choice of the
30 organic water immiscible solvent and the miscible solvent may exert a considerable influence. Examples of water immiscible solvents which may usefully be employed include hexane, chloroform cyclohexane, and toluene. Examples of miscible solvents include acetone, alcohols and acetonitrile. The
35 requirement that the miscible solvent is at least partially

- 4 -

miscible with the water immiscible solvent (and with water) means that the choice of immiscible and miscible solvents are inter-dependent. In the case of crystallisation of solutions of lactose, the preferred solvents are hexane (the immiscible solvent) and acetone (the miscible solvent). The quantities of solvent employed are preferably such as to provide an excess volume of immiscible solvent (typically at least 1.25 and more usually at least 1.5 times the volume of the saturated lactose solution being employed) and a relatively small quantity of the miscible solvent, say no more than 20% by volume being employed.

The solvent mixtures are preferably briskly agitated throughout the period of crystallisation and crystal growth. After the crystal growth phase the particles may be recovered by filtration and are usually washed, e.g. with the miscible solvent to remove excess mother liquor prior to drying. The particles may be subject to further washes, e.g. with ethanol and ethanol/water mixtures to improve the purity. These washes also serve to reduce the quantities of very fine particles present in the product which may be preferable.

The form and size of the crystals may be determined by optical and/or scanning electron microscopy. The rugosity of the particles may be determined by air permeametry which relates the volumetric flow rate (Q) of air through a packed bed of powder compressed to a known porosity to the internal surface area S_o of the powder. The rugosity can then be expressed as the ratio S_o/S_d where S_d is the theoretical surface area (assuming the particles to be spherical). In practice the smoothness of the particles may be readily apparent under the scanning electron microscope and this may render the determination of their rugosity superfluous. Preferably the particles will have a rugosity of no more than 1.5 and most preferably no more than 1.3.

The novel carrier materials are preferably used directly as the sole excipient in dry powder inhalents. However, they may be used in admixture with other excipients although, in general, it is preferred that the excipient comprises at least 80% and

- 5 -

preferably at least 95% by weight of the novel carrier materials of this invention.

05 The novel excipients may be admixed with any suitable pharmacological agent or agents in order to provide a dry powder inhalent composition. Such compositions are believed to be novel and constitute a further aspect of the invention.

10 The average size of the particles of the pharmacological active agent or agents will be such as to facilitate their passage deep into the lower part of the respiratory tract. In general the average particle size should be within the range 0.1 to 10 microns, more preferably 0.5 to 5.0 microns and at least 95% of the particles should have a size within these preferred ranges.

15 The amount of pharmacological agent incorporated into the inhalent composition will generally be from 0.1 to 50% by weight of the composition. The amount will vary with the desired dosage of any particular agent. However, the novel compositions have the advantage that a higher proportion of the pharmacological agent is available to the lower part of the respiratory tract and
20 hence the proportion of any particular agent may be reduced, to one half or even one quarter by weight of the composition compared to a conventional formulation. This increased availability of the active agent also enables agents to be administered by oral inhalation which would not previously have
25 been administered by this route. Thus, agents other than those conventionally employed to treat ailments of the respiratory tract may be administered by this means.

Examples of pharmacological agents which have been administered by oral inhalation include agents with an
30 anti-histamine and anti-allergic action such as sodium cromoglycate and ketotifen, β -agonists, anti-cholinergics such as ipratropium bromide, oxytropium bromide and thiazinamide chloride, sympathomimetic amines such as terbutaline, salbutamol, clenbuterol, pirbuterol, reproterol, procaterol and fenoterol,
35 steroids especially corticosteroids such as beclamethasone

- 6 -

dipropionate, flurisolide budesonide and mucolytics such as ambroxol.

05 Examples of other pharmacological agents which might usefully be incorporated into the novel compositions of this invention include hypnotics, sedatives, tranquillisers, anti-inflammatory agents, anti-histamines, anti-tussives, anti-convulsants, muscle-relaxants, anti-spasmodics, cardiovascular agents, anti-bacterials such as pentamidine, anti-biotics and hypoglycaemic agents.

10 Where appropriate the compositions of this invention may contain a bronchodilator as an additional active agent. The amount of any such bronchodilator will normally not exceed the dosage conventionally employed in its application by inhalation and will preferably be less than is conventionally employed.
15 Examples of useful bronchodilators include isoprenaline, rimiterol, ephedrine, ibuterol, isoetharine, fenoterol, carbuterol, clinbuterol, hexaprenaline, salmifamol, soterenol, trimetoquinol, orciprenaline, terbutaline and salbutamol or a pharmaceutically acceptable salt thereof.

20 The invention finds particular application in the administration of agents which cannot be conveniently administered by other routes. A particular example are peptides such as insulin and growth hormones, ACTH and LHRH analogues.

In addition to the novel carrier and the pharmacologically
25 active agent or agents the compositions of this invention may contain other ingredients such as colouring matter of flavouring agents such as those which are conventionally incorporated into dry powder inhalant compositions. Preferably such ingredients are present in only minor quantities, e.g. less than 10% and more
30 preferably less than 5% by weight of the composition. Such materials will also preferably comprise particles of size comparable with that of the carrier, e.g. 30 to 150 microns.

The compositions may be formulated by dry mixing the active agent and the excipient. The composition may conveniently be
35 encapsulated, e.g. in a hard gelatin capsule suitable for use in

- 7 -

the inhalers which are readily available. The compositions may be formulated into capsules containing a single dose of active material which can be inserted into an appropriate inhaler. Alternatively, they may be placed in a larger container and placed in an inhaler which is designed so as to meter a single dose of the composition into its air passage upon activation. The compositions may be dispensed using any of the conventional inhalers. Their use in dry powder inhalers of all types is strongly preferred. Such inhalers which contain a composition according to this invention are novel and form a further aspect of the invention.

The invention is illustrated by the following examples.

Example 1

Salbutamol sulphate BP was micronised using an air jet mill (Fryma jet mill JM 80) at a pressure of 7.5 bar and a feed rate of 5g/min. The particle size distribution was determined microscopically by measuring the diameter of 500 particles.

Lactose (lactochem Pharmaceuticals), in a size range of 63-90 μm was obtained by sieving (Alpine air jet sieve).

Recrystallised lactose was obtained by crystallisation of the original lactose in a partially miscible mixture of water, hexane and acetone.

Lactose was dissolved in water (2 to 1) in a beaker at 80C. The solution was cooled to room temperature, 75 ml of hexane (Reagent grade) was added to 50ml of the saturated solution and agitated at 500 rpm with a paddle type agitator with four blades, acetone (10ml) (Reagent grade) was then added. The mixture was stirred for 8-12 h, during which time lactose crystals formed. These were washed with acetone, absolute ethanol, 60% ethanol in water and absolute ethanol respectively and dried.

The particle size of the recrystallised lactose was determined with the optical microscope and was found to be in the range of 60-90 μm . The examination of the carrier surface was by scanning electron microscopy. The rugosity of the lactose before and after crystallisation was determined by compressing a mass of

- 8 -

powder equal to its density to a known porosity in the cell of a Fisher Sub-Sieve Sizer. The flow rate through the bed at a fixed pressure differential is transcribed by the instrument to an average particle diameter d_m . The specific surface S_o was calculated from the equation $S_o = \frac{6}{d_m p} \times 10^4$ where p is the powder density. The rugosity before crystallisation was found to be 2.36 whilst the rugosity after recrystallisation was found to be 1.16.

- 10 Samples of drug-lactose blends were prepared in a ratio of 1:67.5 by mixing the micronised drug and the treated lactose with a spatula. The homogeneity of the mixtures was verified by the assay of ten 30 mg samples. The coefficient of variation of the sample content ranged between 1.1-3.0 for the mixtures studied.
- 15 27.4 mg + 1.4 mg of the mixtures containing 400 μ g of salbutamol sulphate was filled into hard gelatin capsules (size 3).

Simulation of patient use

- A diagram of the apparatus is shown in Fig. 1. A powder inhaler device (1) (Rotahaler, Allen & Hanbury's Ltd.) containing an encapsulated dose was assembled in a line conducting dried filtered air at up to 200 l/min. On actuation, the powder was blown into a vertical diffuser (2) 550 mm in length with 2 mm and 70 mm inlet and outlet diameters respectively. Sharp edged conical probes (3) with diameters calculated to give isokinetic sampling were placed at midstream of the diffuser. Air was drawn at 28.3 l/min through a sampler (4) (Anderson 1 CFM Ambient) which comprises a preseparator stage that collects particles with an aerodynamic diameter larger than 10 μ m, and seven separation stages. Stages 0 to 2 have approximate cut-off diameters of 5.5-10 μ m and stages 3 to 7 collect particles less than 5.5 μ m. A final filter trapped particles less than 0.4 μ m.

- Experiments were conducted at air flow rates of 60 and 150 l/min, each using 10 capsules. After deposition, the inhalation device with the capsules, the preseparator, stages 0 to 2, stages 3 to 7 and the filter of the impactor were separately rinsed with

- 9 -

methanol and the washings assayed by HPLC using reversed phase column packed with octadecylsilane (30 cm 3.9 mm i.d.) using 35% 0.013 M ammonium acetate in methanol as the mobile phase and a variable wavelength detector set at 276 nm. The total amount of salbutamol sulphate recovered from each stage was calculated and expressed as a percentage of the total dose discharged.

The mass median diameter of salbutamol sulphate was 2.8 μm with a geometric standard deviation of 1.3.

The results of the effect of surface properties of a carrier on drug deposition are shown in Table I.

Table I

Percentage of drug deposited at various stages using regular lactose and recrystallised lactose.

At air flow rate of 60l/min.

	Regular lactose	Recrystallised lactose
Device	19.7	23.8
Preseparator	57.9	33.6
Stages 0-2	2.8	0.6
Stages 3-7	19.6	42.0

At air flow rate of 150l/min

	Regular lactose	Recrystallised lactose
Device	15.2	24.4
Preseparator	76.8	51.5
Stages 0-2	2.6	2.6
Stages 3-7	5.4	22.0

- 10 -

Example 2

A double blind randomised cross-over trial was carried out to compare the effects of a commercial formulation comprising salbutamol sulphate and a conventional lactose carrier with a composition according to this invention containing the same proportions of salbutamol sulphate and a modified lactose of this invention prepared in the manner described in Example 1. Eleven moderate to severe stable atopic asthmatic patients took part in the trial (FEV₁ <80% predicted; >15% reversibility. FEV₁ is Forced Expiratory Volume in 1 second). The trial was carried out using conventional dry powder inhalers. The commercial formulation produced a mean increase in FEV₁ of 21.4%. The formulation according to this invention produced a mean increase in FEV₁ of 27.5%. The difference 6.1% was significant (paired t-test; p <0.05; confidence interval 0.64-11.52).

- 11 -

What we claim is

1. A particulate carrier suitable for use in dry powder inhaler compositions having an average particle size of from 5.0 to 1000 microns and a rugosity of less than 1.75.
 - 05 2. A carrier according to Claim 1 characterised in that the carrier is a particulate crystalline sugar.
 3. A carrier according to Claim 2 characterised in that the crystalline sugar is selected from the group comprising glucose, fructose, mannitol, sucrose and lactose.
 - 10 4. A carrier according to Claim 3 characterised in that the crystalline sugar is lactose.
 5. A carrier according to any of Claim 1 to 4 characterised in that the particles have an average particle size of from 30 to 250 microns.
 - 15 6. A carrier according to any of the preceding claims characterised in that the particles have a rugosity of not more than 1.5.
-
7. A dry powder inhalent composition which comprises a excipient in admixture with at least one pharmacological agent which is
 - 20 characterised in that the excipient comprises a particulate carrier according to any of Claims 1 to 6.
 8. A composition according to Claim 7 characterised in that the excipient comprises at least 80% by weight of a particulate carrier according to any of Claims 1 to 6.
 - 25 9. A composition according to Claim 8 characterised in that the excipient consists essentially of a particulate carrier according to any of Claims 1 to 6.
 10. A composition according to any of Claims 7 to 9 characterised in that it comprises from 0.1 to 50% by weight of a
 - 30 pharmacological agent.
 11. A composition according to any of Claims 6 to 10 characterised in that the pharmacological agent is a particulate solid having an average particle size of from 0.1 to 10.0 microns.

- 12 -

12. A composition according to any of Claims 6 to 11 characterised in that the pharmacological agent is a β agonist, a steroid or sodium chromoglycate.
- 05 13. A composition according to any of Claims 6 to 11 characterised in that the pharmacological agent is a peptide.
14. A composition according to Claim 13 characterised in that the pharmacological agent is selected from the group comprising insulin and growth hormones such as ACTH and LNRH analogues.
- 10 15. A composition according to any of claims 6 to 11 characterised in that the pharmacological agent is an anti-bacterial agent.
16. A composition according to claim 15 characterised in that the anti-bacterial agent is pentamidine.
- 15 17. A process for the production of a particulate carrier according to any of Claims 1 to 6 which comprises precipitating the carrier from a saturated aqueous solution by the addition of at least an equal volume of a water immiscible organic solvent and a quantity of a solvent which is miscible with water and with the water immiscible solvent which is at least 5% by volume of
- 20 the volume of the aqueous solution.
18. A process according to Claim 16 characterised in that the water immiscible solvent is selected from the group comprising hexane, chloroform cyclohexane and toluene.
- 25 19. A process according to either of Claims 17 or 18 characterised in that the water miscible solvent is selected from the group comprising acetone, ethanol propanol or butanol and acetonitrile.
20. An encapsulated dry powder inhalent composition according to any of claims 7 to 14.
- 30 21. A dry powder inhaler characterised in that it contains a dry powder inhalent composition according to any of claims 7 to 14 or 20.
22. A carrier according to Claim 1 substantially as hereinbefore described with reference to the foregoing examples.

1 / 1




Fig. 1

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 91/00103

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶

According to International Patent Classification (IPC) or to both National Classification and IPC

IPC⁵: A 61 K 9/72, A 61 K 47/26

II. FIELDS SEARCHED

Minimum Documentation Searched ⁷

Classification System

Classification Symbols

IPC⁵

A 61 K

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched ⁸

III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹

Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	DE, A, 2851489 (FISONS) 31 May 1979 see claims 1-4,9,15; page 4	1-5,7-12,22
Y	---	17,19
X	WO, A, 8705213 (CHIESI FARMACEUTICI) 11 September 1987 see claims 1,2,3,8,9,11,12,13,14	1-5,7-12, 15,22
Y	---	17,19
Y	DE, C, 292149 (FARBWERKE MEISTER LUCIUS & BRUNING) 24 May 1916 see claims; page 1, column 1, paragraph 1	17,19

¹⁰ Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search

15th April 1991

Date of Mailing of this International Search Report

22.05.91

International Searching Authority

EUROPEAN PATENT OFFICE

Signature of Authorized Officer

Mme. M. van der Drift

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

GB 9100103

SA 43670

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 14/05/91
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE-A- 2851489	31-05-79	GB-A- 1571629	16-07-80
		AU-B- 525470	11-11-82
		AU-A- 4210078	07-06-79
		BE-A- 872319	28-05-79
		FR-A, B 2423218	16-11-79
		JP-A- 54084022	04-07-79
		US-A- 4199578	22-04-80
WO-A- 8705213	11-09-87	AU-B- 597964	14-06-90
		EP-A, B 0239798	07-10-87
		EP-A- 0258356	09-03-88
		JP-T- 63502895	27-10-88
		ZA-A- 8701523	24-08-87
DE-C- 292149		None	

THIS PAGE BLANK (USPTO)